

Aneurysms & Dissections

↳ Dilatation

↳ Rupture of
the Artery

Dr. Bushra Al-Tarawneh, MD

Anatomical pathology

Mutah University

School of Medicine-

Department of Microbiology & Pathology
lectures 2022




Aneurysms

← A congenital or acquired dilations of blood vessels or the heart, could be:

1. **“True”** : all three layers of the artery (intima, media, & adventitia) or the wall of the heart; e.g. atherosclerotic, congenital vascular aneurysms, ventricular aneurysms after MI


1. **“false”** : a wall defect leads to the formation of an extravascular hematoma that communicates with the intravascular space (“pulsating hematoma”)



Aneurysms – Types by shape

Risk factors
 Risk factors - الأسباب
 shape - الشكل
 locations - المواقع
 false aneurysm - متزيف

- ← **Saccular aneurysms:** *→ Inside is involved* discrete outpouchings ranging (5-20 cm) in diameter, often with a contained thrombus.
- ← **Fusiform aneurysms:** *→ Both sides are affected* circumferential dilations up to 20 cm in diameter, most commonly involve aortic arch, abdominal aorta, or iliac arteries.



Aneurysms – Pathogenesis

Alterations in **SMCs** or **ECM** → compromise **structural integrity** of the arterial media

1. **Inadequate or abnormal connective tissue synthesis:**

← **Marfan syndrome:** defective synthesis of scaffolding protein **fibrillin** → ↑**TGF-β** activity → weak of elastic tissue → dilation in the aorta.

Male
tall, protruding jaw
long bones
↓
defective Fibrillin

← **Ehlers-Danlos syndrome IV:** Defective type III collagen synthesis → weak vessels → aneurysm formation.

↓
couldn't
w/stand
the pressure

Aneurysms – Pathogenesis

2. The balance of collagen degradation and synthesis is altered by inflammation and associated proteases.

Increased matrix metalloprotease expression by macrophages in atherosclerotic plaque → aneurysm by degrading arterial ECM.

↳ Because of the Inflammation caused by the plaque → Component of BV is altered → weakness of BV wall.

3. The vascular wall is weakened through loss of smooth muscle cells or the synthesis of noncollagenous or nonelastic extracellular matrix.

Aneurysms – Pathogenesis


- ← **Medial ischemia** may lead to “degenerative changes” of the **aorta**; Ischemia → **smooth muscle cell loss** → **scarring** and **loss of elastic fibers** → **inadequate extracellular matrix synthesis** → **production** of increasing amounts of amorphous ground substance (**glycosaminoglycan**).
- ← Histologically, these changes recognized as **cystic medial degeneration**

Aneurysms – cystic medial degeneration

Degenerated

Elastic stain

Normal





Abdominal Aortic Aneurysm (AAA) ^{"triple A"}

- ← Aneurysms occurring as a consequence of **atherosclerosis** form most commonly in **abdominal aorta & common iliac arteries**. *Associated w/ it.*
- ← More frequently in men & in smokers & rarely before 50.
- ← Atherosclerotic plaques → compromise the **diffusion** of nutrients & wastes between vascular lumen & arterial wall → deleterious effects on SMCs.
- ← Also atherosclerotic lesions → Inflammatory infiltrates → release **proteolytic enzymes** → ECM degradation
- ← Combination of these, the media undergoes degeneration & necrosis → arterial wall thinning → dilation

AAA- Morphology

- ← AAAs typically occur between the renal arteries & the aortic bifurcation; can be saccular or fusiform & up to 15 cm in diameter and 25 cm in length.
- ← In the vast majority extensive atherosclerosis is present, with thinning & focal destruction of the underlying media.
- ← The aneurysm sac usually contains bland, laminated, poorly organized mural thrombus ...can fill much of the dilated segment.
↳ Inside the lamination
- ← Not infrequently, AAAs are accompanied by smaller iliac artery aneurysms.

AAA- Morphology



AAA- Clinical Manifestations

- ← Most AAA are asymptomatic & discovered incidentally as **An abdominal mass** (often palpably pulsating) that simulates a tumor.
- ← **Obstruction** of a vessel branching off the aorta.
- ← **Embolism.**
- ← **Impingement on adjacent structures.**
- ← **Rupture** into the peritoneal cavity or retroperitoneal tissues → massive, often fatal hemorrhage.

↳ Because it's
a main Artery
w/ ↑↑ Blood flow

AAA- Clinical Consequences

- ← The risk for rupture is related to the size of AAAs. (4 cm in diameter or less almost never burst, 4-5 cm do so at a rate of 1% per year, 11% per year for AAAs 5-6 cm, & 25% per year for aneurysms >6 cm in diameter.

more than 5cm → SURGERY! Associated w/ rupture

- ← Aneurysms 5 cm in diameter or larger are managed surgically.
- ← Timely intervention is critical; (mortality in elective procedures ~ 5%, whereas in emergency surgery after rupture ~ 50%)

Thoracic Aortic Aneurysm

Risk factors &

Most commonly associated with hypertension, bicuspid aortic valves, & Marfan syndrome & Manifest with the following signs & symptoms:

- Respiratory or feeding difficulties ⇒ *In children*
- Persistent cough from irritation of the recurrent laryngeal nerves.
- Pain caused by erosion of bone.
- Cardiac disease due to valvular insufficiency, narrowing of the coronary ostia, or aortic valvular incompetence.
- Aortic dissection or rupture.

Aortic Dissection

- ← Aortic dissection occurs when blood separates the laminar planes of the media to form a blood-filled channel within the aortic wall. ⇒ The blood is collected w/ in the media
- ← Catastrophic if the dissection ruptures through the adventitia & hemorrhages into adjacent spaces.
- ← two groups of patients:
 - (1) Men **40-60 years** with antecedent **hypertension** (>90% of cases). (EPO)
 - (2) **Younger adults** with systemic or localized abnormalities of connective tissue affecting the aorta (e.g., **Marfan syndrome**).
or EDS

Aortic Dissection - Pathogenesis

- ← Hypertension is the major risk factor for aortic dissection.
- ← Narrowing of the vasa vasorum → diminished flow through vasa vasorum → degenerative changes in ECM & variable loss of medial SMCs.
- ← Abrupt, transient increase in blood pressure, as may occur with cocaine abuse, is also known to cause aortic dissection.
- ← The trigger for the intimal tear is not known in most cases. Nevertheless, once the tear has occurred, blood under systemic pressure dissects through the media along laminar planes.


Aortic Dissection - Morphology

16



mostly the intimal tear marking origin point is found in the ascending aorta within 10 cm of the valve. Dissection plane can extend retrograde toward the

wt in the Media of BV.



Aortic Dissection - Morphology

Associated w/ Hypertension

Dissection plan usually lies between the middle and outer thirds of the media.



Aortic Dissection – Clinical presentation

18

- ← The classic clinical symptom of aortic dissection is the **sudden onset of excruciating tearing or stabbing pain** in the anterior chest, radiating to the back between the scapulae, & moving downward as the dissection progresses.
- ← The most common cause of death is rupture of the dissection into the pericardial, pleural, or peritoneal cavity.
- ← Common clinical presentations stemming from cardiac involvement include tamponade, aortic insufficiency, and myocardial infarction.

Aortic Dissection – Clinical presentation

← Aortic dissections generally are classified into two types

Proximal lesions (type A dissections), involving the ascending aorta with or without involvement of the descending aorta (DeBakey type I or II) respectively)...

Needs rapid diagnosis & institution of intensive anti-hypertensive therapy coupled with surgery.. **Worse outcome** (most common)

Distal lesions (type B dissections), beginning beyond the subclavian artery (DeBakey type III) ... Mostly can be managed conservatively.. **Better outcome**; 75% survival rate whether they are treated with surgery or with antihypertensive only.

Stanford

DeBakey

type A

type I



type II



type B

type III



should involve
the Ascending
Aorta

Veins and Lymphatics

90% of clinical venous disease caused by Varicose veins
and phlebothrombosis/thrombophlebitis

Varicose veins ⇒ Prominent vascular lesions in the leg.

- ← Abnormally **dilated tortuous** veins produced by chronically (1) increased intraluminal pressures & (2) weakened vessel wall support.
- ← Venous valves incompetent → lower-extremity stasis, congestion, edema, pain, & thrombosis.
- ← **Locations:** typically, superficial veins of the upper & lower leg.
- ← **Risk factors:** Obesity, female sex, pregnancy, & familial tendency.
- ← **Clinical features:** persistent edema & secondary ischemic, skin changes, including stasis dermatitis and ulcerations.

Varicose veins– Clinical presentation

* Mostly Associated w/ long time of- standing



(الفرق بينهم في الـ)

24

Thrombophlebitis & Phlebothrombosis

- ← Two terms are largely interchangeable designations for **venous thrombosis** accompanied by **inflammation**.
- ← 90% of cases are due to thrombosis of deep leg veins (DVT)
- ← **Risk factors for DVT:** **Prolonged immobilization**, postoperative state, congestive heart failure, pregnancy, oral contraceptive use, malignancy, obesity, male sex, & age over 50 year. *Female > Male*
- ← **Clinical manifestation:** few reliable signs or symptoms; distal edema, cyanosis, superficial vein dilation, heat, tenderness, redness, swelling, & pain. However, many DVTs are asymptomatic, and the absence of findings does not exclude their presence.

⊕ Unilateral swelling

DVT ← *أحد أعراضها*

Thrombophlebitis & Phlebothrombosis

← Pulmonary embolism is a common & serious clinical complication of DVT, resulting from fragmentation or detachment of the venous thrombus.

VE ← In many cases, the first manifestation of thrombophlebitis is a pulmonary embolus.

← Depending on the size and number of emboli, the outcome can range from resolution with no symptoms to death

