The background features a stylized autumn landscape. On the left, a red brick barn with a yellow bell is visible. A yellow school bus is driving on a winding road that curves across the scene. The foreground is filled with various autumn trees in shades of orange, yellow, and brown. In the bottom right corner, there are several pumpkins. The sky is light blue with a few scattered autumn leaves.

Sexually transmitted infections (Public Health Perspective)

Dr. Israa Al-Rawashdeh MD,MPH,PhD
Faculty of Medicine
Mutah University
2023



Definition:

- Sexually Transmitted Infection (STI): are infections that result from the transmission of a pathogenic organism predominantly through sexual contact.
- Sexually Transmitted Disease (STD): refers to a recognizable disease state that has developed from an infection.



- Human beings are the only natural reservoir for STI etiological agents.
- A person can have an STI without having obvious symptoms of disease (asymptomatic)
- Several STI can occur in one same individual.




Albrecht Durer's (1471-1528) woodcut of a syphilitic man covered in chancres.

Sexually Transmitted Infections

Magnitude:

- STIs have a great impact on sexual and reproductive health worldwide.
 - More than 1 million STIs are acquired every day.
 - There are approximately 36.7 million people living with HIV by the end of 2015 , 2.1 million of which were newly infected.
 - More than 500 million people are living with genital HSV (herpes) infection.
 - In 2016, WHO estimated 376 million new infections with 1 of 4 STIs among people 15–49 years of age :
 - Chlamydia (127 million),
 - Gonorrhoea (87 million),
 - Syphilis (6.3 million), and
 - Trichomoniasis (156 million).



Risk factors


Factors that may increase STIs' risk include:

- **Age** (Half the new STIs occur in people between the ages of 15 and 24)
- **Unprotected sexual activities.**
- **Having sexual contact with multiple partners.**
- **Previous history of STIs.**
- **Misuse of alcohol or use of recreational drugs.** Increase susceptibility to participate in risky behaviors.
- **Injecting drugs.** Needle sharing spreads many serious infections, including HIV, hepatitis B and hepatitis C.
- **Transmission from mothers to infants** Certain STIs — such as gonorrhea, chlamydia, HIV and syphilis — can be passed from mothers to their infants during pregnancy or delivery.
- **Immune status of the individual**



The rate of spread (R_0) of STIs in populations: **infectivity (β)**, **rate of exposure between infected and susceptible individuals (C)**, and **duration of infection (D)**, and the factors that affect those variables.

Framework:





Sexually Transmitted Infections

- More than 30 different bacteria, viruses and parasites are known to be transmitted through sexual contact.
 - Eight of these pathogens are responsible for the greatest incidence of STIs.
 - Of these 8 infections; 4 are currently **curable**: syphilis, gonorrhoea, chlamydia and trichomoniasis.
 - The other 4 are viral infections which are **incurable**: hepatitis B, herpes simplex virus (HSV or herpes), HIV, and human papillomavirus (HPV).
 - Symptoms or disease due to the incurable viral infections can be reduced or modified through long term treatment.
- 
- 

STIs: Pathogens

Bacteria:

Gonorrhea (*Neisseria gonorrhoeae*)

Chlamydia (*Chlamydia trachomatis*)

Syphilis (*Treponema pallidum*)

Chancroid (*Haemophilus ducreyi*)

Viruses:

Genital warts and cervical—mainly-cancer (human papillomavirus)

Genital herpes (herpes simplex virus)

Hepatitis B (hepatitis B virus)

AIDS (HIV)


Parasites and protozoa:

Trichomoniasis (*Trichomonas vaginalis*)

Pubic lice (*Phthirus pubis*)

Candidiasis or "yeast infection"??

Since the 80's: HIV, the new, devastating, STI



Activate Windows
Go to Settings to activate W



Sexually Transmitted Infections

- 300 million women have an HPV infection, the primary cause of cervical cancer.
- An estimated 240 million people are living with chronic hepatitis B globally.

Both HPV and hepatitis B infections are preventable with vaccination.



Sexually Transmitted Infections: Impact

- Impact: STIs can have serious consequences beyond the immediate impact of the infection itself.
 - ❑ **HIV risk** – the presence of a sexually transmitted infection, such as syphilis, gonorrhoea, or HSV, increases the risk of acquiring or transmitting HIV infection (by two to three times, in some populations)
- 

Sexually Transmitted Infections

❑ Mother-to-child transmission of STIs can result in **Fetal and neonatal impact:**

- Stillbirth, neonatal death,
- Low-birth-weight and prematurity,
- Sepsis, pneumonia, neonatal conjunctivitis, and congenital deformities.

Syphilis in pregnancy leads to over 300 000 fetal and neonatal deaths each year, and places an additional 215 000 infants at increased risk of early death

Worldwide, up to 4000 newborn babies become blind every year because of eye infections attributable to untreated maternal gonococcal and chlamydial infections.




Sexually Transmitted Infections

- Impact:

- **Cervical cancer** – the human papillomavirus HPV (16 and 18) infection is responsible for an estimated 530 000 cases of cervical cancer and 264 000 cervical cancer deaths each year

- STIs such as gonorrhoea and chlamydia are major causes of **pelvic inflammatory disease** (PID) and **infertility** in women.



Sexually Transmitted Infections

- Sign and symptoms of STIs:

Male:

- Urethral discharge
- Burning and pain during urination

Female:

- Abnormal vaginal discharge
- Burning and/or increased frequency of urination
- Lower abdominal pain

Both:

Pain, itch, Papules, vesicles, erosion/ulcers or fleshy growths in and around genitalia, perineum and anus/rectum, oral cavity and occasionally on other sites.

Swelling in inguinal, anal region or scrotum.



Prevention and control:

- Prevention and control of STIs are based on the following five major strategies :
 1. **Accurate risk assessment and education and counseling** of persons at risk regarding ways to avoid STIs through changes in sexual behaviors and use of recommended prevention services
 2. **Pre-exposure vaccination** for vaccine-preventable STIs
 3. **Identification** of persons with an asymptomatic infection and persons with symptoms associated with an STI
 4. Effective **diagnosis, treatment, counseling, and follow-up** of persons who are infected with an STI
 5. Evaluation, treatment, and counseling of **sex partners** of persons who are infected with an STI
- 
- 
- 
- 

Primary Prevention Methods

1. Pre-Exposure Vaccination:

Pre-exposure vaccination is one of the most effective methods for preventing transmission of HPV, HAV, and HBV.

HPV vaccination is recommended routinely for males and females aged 11 or 12 years and can be administered beginning at age 9 years.

Hepatitis B vaccination is recommended for all unvaccinated, uninfected persons who are sexually active with more than one partner or are being evaluated or treated for an STI. hepatitis B vaccines are recommended for MSM, persons who inject drugs, persons with chronic liver disease, and persons with HIV or hepatitis C infections who have not had hepatitis B.

TABLE 2**HPV Vaccine Dosing Schedules Based on Age**

Age (males and females)	Doses	Schedule
9-14 y*	2-dose series [†]	Dose 1: 0 mo Dose 2: 6-12 mo
15-26 y	3-dose series	Dose 1: 0 mo Dose 2: 1-2 mo Dose 3: 6 mo

three-dose schedule also applies for immunocompromised persons initiating vaccination at ages 9 through 26 years; and to adults initiating vaccination at ages 27 through 45 years.

Source: Meites et al. *MMWR Morb Mortal Wkly Rep.* 2016.⁷



International Hepatitis B Vaccine Schedules For All Ages

Vaccine	Dose 1	Dose 2	Dose 3	Dose 4
 3-dose vaccine series for infants < 1 year	Within 24 hours of birth ✓	1 month after dose 1 (At least 4 weeks of age) ✓	6 months after dose 1 (At least 24 weeks of age) ✓	
 3-dose vaccine series for children ≥ 1 year and adults	Now ✓	1 month after dose 1 ✓	6 months after dose 1 ✓	

Primary Prevention Methods


2. Barrier methods

- When used correctly and consistently, **condoms** offer one of the most effective methods of protection against STIs, including HIV.
- Female condoms are effective and safe, but are not used as widely by national programmes as male condoms.

Contraceptive methods that are not mechanical barriers offer no protection against HIV or other STIs

3. Male Circumcision

- Male circumcision reduces the risk for HIV infection by 50%–60% and certain STIs including high-risk genital HPV infection and genital herpes among heterosexual men.



Sexually Transmitted Infections

4. Counselling and behavioural approaches

- Comprehensive sexuality education, STI and HIV pre- and post-test counselling;
- Safer sexual behaviours/risk-reduction counselling, condom promotion;
- STI interventions targeted to key populations, such as sex workers, men who have sex with men (MSM) and people who inject drugs;
- Improve people's ability to **recognize** the symptoms of STIs and increase the likelihood they will **seek care** or encourage a sexual partner to do so.
- Abstinence and Reduction of Number of Sex Partners Abstinence from oral, vaginal, and anal sex and participating in a long-term, mutually monogamous relationship (Commitment) with a partner known to be uninfected are prevention approaches to avoid transmission of STIs.

Sexually Transmitted Infections

5. Pre and Post- exposure Prophylaxis for STIs and HIV.

- Example:
- Doxycycline prophylaxis has been examined for preventing bacterial STIs
- Antiviral drugs: fixed-dose combination of emtricitabine (FTC) and either tenofovir disoproxil fumarate (TDF) or tenofovir alafenamide (TAF)

Sexually Transmitted Infections

- **Unfortunately,**
 - Lack of public awareness,
 - Lack of training of health workers,
 - Long-standing, widespread **stigma** around STIs remain barriers to greater effective use of these interventions.





Sexually Transmitted Infections

- **Management:**
- **Case management:**
 - STI case management is the care of a person with an STI-related syndrome or with a positive test for one or more STI.
- The components of case management include:
 - History taking, examination, correct diagnosis, early and effective treatment,
 - Advice on sexual behaviour, promotion and/or provision of condoms,
 - Partner notification and treatment,
 - Case reporting and
 - Clinical follow-up as appropriate

Treatment of STIs

- Effective treatment is currently available for several STIs.
- Three bacterial STIs (chlamydia, gonorrhoea and syphilis) and one parasitic STI (trichomoniasis) are generally curable with existing, effective **single dose regimens of antibiotics**.
- For herpes and HIV, the most effective medications available are **antivirals** that can modulate the course of the disease, though they cannot cure the disease.
- For hepatitis B, **immune system modulators (interferon)** and antiviral medications can help to fight the virus and slow damage to the liver.
- **Resistance of STIs – in particular gonorrhoea** – to antibiotics has increased rapidly in recent years and has reduced treatment options.
- The emergence of decreased susceptibility of gonorrhoea to the “last line” treatment option (oral and injectable cephalosporins) + antimicrobial resistance already shown to penicillins, sulphonamides, tetracyclines, quinolones and macrolides make gonorrhoea a multidrug-resistant organism.

Sexually Transmitted Infections



• Management:


- The identification of consistent groups of symptoms and easily recognized signs (syndromes) to guide treatment, without the use of laboratory tests.

This is called **syndromic management**.

- This approach mainly followed (In poor countries), allows health workers to diagnose a specific infection on the basis of observed syndromes as:
 - vaginal discharge,
 - urethral discharge,
 - genital ulcers,
 - abdominal pain



STI/RTI SYNDROMIC CASE MANAGEMENT





Sexually Transmitted Infections



- 
- **Management:**
 - **STI case management**
 - Syndromic management is simple, assures rapid, same-day treatment, and avoids expensive or unavailable diagnostic tests for patients that present with symptoms.
 - This approach results to **overtreatment** and **missed** treatment as majority of STIs are asymptomatic.
 - Therefore, in addition to syndromic management, **screening** strategies are essential.
 - To interrupt transmission of infection and prevent re-infection, treating **sexual partners** is an important component of STI case management.
- 