

Enzymology- An overview-1

Introduction about it.


* Turn over number \Rightarrow
is the number of substrate molecules
to be convert into Product molecules
Per unit enzyme Per one second.

* If the enzyme
is found in the dead
animal under the land
the Petroleum will produce
very fast. than if the
enzyme not found.

under the high
temperature and
pressure.

البتروول = منتج بعد
منه في اقل وقت
زهرًا له لم وجود
انزيمات.

Enzymes- An introduction



- Biologic (organic catalysts) polymers that catalyze the chemical reactions.

Another catalyst →
 → (inorganic) like metal.

acceleration of the reaction } ⇒ acceleration

لا يمتص ولا يتغير كل تلك بعد التحويل

- Enzymes are neither consumed nor permanently altered as a consequence of their participation in a reaction.

* The enzyme when it enter to the reaction will exit like it was before it enter.

hydrogenation →
 درجة من سرعة (الحمأة)
 حمض و جين للرابطة - المتناحية

- With the exception of catalytic RNA molecules, or ribozymes, enzymes are proteins.

they have catalytic activity → make "processing of the RNA" or modification.

while the inorganic catalysts these will be convert about (10^9) in one second.

(not Proteins) short stretches from (RNA) (90-300) converting immature (RNA) (inactive) into mature (RNA) by post-translational modification

- In addition to being highly efficient, enzymes are also extremely selective catalysts.

is the number of substrate molecules to be converted by one molecule of Per time unit. the enzyme can be work on (10^6-10^8)

- Thermolabile, site specific, with a high turn over number compared to the inorganic catalysts.

يتأثر بالحرارة كل انزيم موجود في منطقة القابلة

جودتها كجودتها عالية
 جودتها لينة + جودتها جودتها

cofactors

20 characteristics of enzymes

تتفرقة حيث للبروتين الكروي والليفاني
 كروي 107
 ليفي 102
 ← الأول
 ← الثاني
 * axial ratio :

* غالبية الإنزيمات تكون
 هيئاتها كروية لأنها تكون
 مكونة من جودات كروية.
 * وإغلبها تحت دقات
 اثنائي بسبب تركيبها.


Enzymes are both intracellular and extracellular catalysts

Some are globular proteins and few are RNA-based molecules

Some enzymes need coenzymes or cofactors

Enhance the speed of biochemical reactions

Forms enzyme-substrate complex



Lowers the activation energy

Active site contains less hydrophobic amino acids

Produces product using specific substrate

Sensitive to temperature, pH, and substrate concentration

Required in very less amount compared to chemical catalyst

Active site contains 3 to 12 amino acids

The enzyme is convert to the substrate after it

Enzymes can be recycled or reused

Enzymes are larger than substrate

Function can be inhibited by inhibitors

التسمية العلمية
تحتوي على نوع التفاعل
والمادة المتفاعلة

Nomenclature of enzymes

-In most cases, enzyme names end in **-ase**

urea
→ uricase

-The common name for a hydrolase is derived from the substrate

Urea: remove -a, replace with **-ase** = urease

Lactose: remove -**ose**, replace with **-ase** = lactase

- Other enzymes are named for the substrate and the reaction catalyzed

✓ Lactate dehydrogenase $-H_2$

✓ Pyruvate decarboxylase $-CO_2$

- Some names are historical - no direct relationship to substrate or reaction type

Catalase

Pepsin → in the stomach

بوالبنكرياس

Chymotrypsin → breakdown of peptide bond
تفكيك الروتين الى امينو اسيد

Trypsin

Peptidase } ⇒ hydrolases

Classification of Enzymes

- Enzyme Commission (EC) – according to International Union of Biochemistry and Molecular Biology (IUBMB)

- Each enzyme was given 4 digit numbers ⁶ [1.2.3.4]

1st one of the 6 major classes of enzyme activity
^{CNC}
^{⇒ (Commission numerical code)}

^{① oxidation}

2nd the subclass (type of substrate or bond cleaved)

3rd the sub-subclass (group acted upon, cofactor required, etc...)


4th a serial number... (order in which enzyme was added to list)

1- Oxidoreductases (EC.1) catalyze redox reactions, such as
redox reaction (Alcohol dehydrogenase [EC 1.1.1.1])

- Reductases
- Oxidases

متراجات
 وبن اذا كان في H^+ زيادة
 زيادة نصفي الوتق سر جلي زيادة

تتأثر ال pH بسبب فقد (H^+)
يعني \uparrow pH




2- Transferases (EC.2) transfer a group from one molecule to another,
 such as (Hexokinase [EC 2.7.1.2])

- Transaminases catalyze transfer of an amino group
- Kinases transfer a phosphate group *ATP/ADP*


All kinase is a Transferases.

from (S-adenosine methionine) SAM
will be converted to (SAH)
→ (S-adenosine Homocysteine)



5- Isomerases (EC.5) catalyze intramolecular rearrangements, such as
 (Alanine racemase [EC 5.1.1.1])


- ✓- Epimerases
- ✓- Mutases




6- Ligases (EC.6) catalyze a reaction in which a C-C, C-S, C-O, or C-N bond is made or broken, such as
 (Isoleucine-tRNA ligase [EC 6.1.1.5])

Binding bonds

need energy for reactions



This type only require energy for completing the reaction.





Active site

- Takes the form of a cleft or pocket
- Takes up a relatively small part of the total volume of an enzyme
- Substrates are bound to enzymes by multiple weak attractions
- The specificity of binding depends on the precisely defined arrangement of atoms in an active site
- The active sites of multimeric enzymes are located at the interface between subunits and recruit residues from more than one monomer

→ It has a particular amino acids which called highly reactive groups. Like :- ① hidden ② glutamic acid

رصيدا لاصية بقية على ترتيب الذرات



more than one protein molecule the active site will be in between the peptide bond.



* The enzyme should be 3D to be complementary to the shape of substrate.

Types of groups forming the active site :-

- ① Hydroxy - amino acid containing
- ② Acidic Amino
- ③ Basic Amino
- ④ imidazole



The enzyme should have a 3D structure form to complete its function.


Enzyme substrate binding

-Two models have been proposed to explain how an enzyme binds its substrate: the **lock-and-key model** and the **induced-fit model**.

✗ If the substrate is not super malleable in the active site there will be no reaction.


- **Lock-and-Key Model of Enzyme-Substrate Binding**, in this model, the active site of the unbound enzyme is **complementary in shape** to the substrate.

- "lock and key model" **accounted for the exquisite specificity of enzyme-substrate interactions**, the **implied rigidity** of the enzyme's active site failed to account for the dynamic changes that accompany catalysis.



Induced-Fit Model of Enzyme-Substrate Binding

- In this model, the enzyme changes shape on substrate binding.
- The active site forms a shape complementary to the substrate only after the substrate has been bound.
- When a substrate approaches and binds to an enzyme they induce a (conformational) change, a change analogous to placing a hand (substrate) into a glove (enzyme).




* The changes will be after the substrate binding to the enzyme.

* This mechanism is complete with hormones and other substances.

Mechanism of Action of Enzymes

- Enzymes are catalysts and increase the speed of a chemical reaction without themselves undergoing any permanent chemical change. They are neither used up in the reaction nor do they appear as reaction products.
- The basic enzymatic reaction can be represented as follows:



- Where E represents the enzyme catalyzing the reaction, S the substrate, the substance being changed, and P the product of the reaction.
- The mechanism of action of enzymes can be explained by two perspectives:
 - 1- Thermodynamic changes
 - 2- Processes at the active site


Thermodynamic changes ①

- All enzymes accelerate reaction rates by providing transition states with a lowered ΔG^\ddagger for formation of the transition states.
- The lower activation energy means that more molecules have the required energy to reach the transition state.

* The enzyme allow the reaction complete with the least activation of energy.
 To accelerate the reaction.

* The bond between the functional group in the active site and the substrate must be

- ① hydrogen bond
- ② hydrophobic interaction
- ③ disulfide bond
- ④ Van der Waals forces / electrostatic



Processes at the active site (2)

1- **Catalysis by proximity:** for the molecules to react they

① must come within bond-forming distance of one another.

When an enzyme binds substrate molecules at its active

② site, it creates a region of high local substrate

concentration. *Should be crowdedness between the substrate molecules.*

③ Enzyme-substrate interactions orient reactive groups and bring them into proximity with one another.

orientation of the reactive groups in the active site - not in the binding site of the substrate.

لا يحد على الروابط ان تكون قوية بين الاضراس والماند المقابل

2- **Acid base catalysis:** the ionizable functional groups of aminoacyl

side chains of prosthetic groups contribute to catalysis by acting as

(acids or bases). *Acidic amino acid like ① glutamic acid / ② Aspartic acid. Basic amino acid like ① histidine ② Asn ③ Gln* } occur the ionization for these groups } (-ive) for Acidic } (+ive) for Basic }

- General acid catalysis involves partial proton transfer from a donor to lower the free energy of the transition state.

- General base catalysis involves partial proton abstraction from an acceptor to lower the free energy of the transition state.

الاضراس سهل العكس بيت العزيمتات لذلك سهل المقابل

1] General Acid catalysis :-

Partial Proton transfer from a donor to lower the activation energy.

* To lower the activation energy should the ionization will be occur.

} Donor the
(H^+)

2] General Basic catalysis :-

Is the same with the Acidic

but involves Partial Proton abstraction from the acceptor to arrive the transition state.

} accept the
(H^+)

→ Is also called lytic enzymes also they involves All digistic enzymes.

3- **Catalysis by strain:** ^{ليس} enzymes that catalyze the lytic ^{not for All enzyme} reactions involve breaking a covalent bond typically bind their substrates in a **configuration slightly unfavorable** ^{ما تكافؤ} for the bond that will undergo **cleavage.** ^{غير مريحة / محبة} binding / غير مريحة

^{تقسيم} ^{الانزيم} ^{ترتبط بشكل مائل و ضعيف كل الانزيم}

4- **Covalent catalysis:** ^{for not All enzyme} accelerates reaction rates through **transient** formation of enzyme-substrate covalent bond.

Three stages in covalent catalysis:

1- Nucleophilic reaction ^{أخذ ورد البروتونات} between enzyme and substrate

2- Electrophilic withdrawal ^{إفقد ورد الإلكترونات} of electrons from substrate

3- Elimination reaction (reverse of stage 1) ^{regulation the ionization.}

* If the ionization state is increase the bonds between the molecules will be breaking down.

* the ionization in reactive groups should be under regulation.

not for all enzymes

بعد الاستعمال (Cu / Fe)

5- Metal Ion catalysis

- Two classes of metal ion dependent enzymes:

1- **Metalloenzymes** contain tightly bound transition metal ions

(Fe²⁺, Fe³⁺, Cu²⁺, Zn²⁺, Mn²⁺)

عبر مطلوب بالضرورة } اذا كان غير موجود يحتاجه لكن بشكل ضئيف

2- **Metal-activated enzymes** loosely bind metal ions (alkali or alkaline metal including Na⁺, K⁺, Mg²⁺ and Ca²⁺)

- Metal ions enhance catalysis in three major ways:

1- Binding to and orienting substrates for reaction as Mg²⁺ binding to ATP

① in between them (enz and sub) | ② pushed them
② pushed them (enz to the sub) | (sub to enz)

2- Mediating **redox reaction** through changes in oxidation state such as reduction of O₂ to H₂O through electron transfer

the increasing the electrons will increasing the ionization to be breakdown down the interaction between the binding molecules.

3- **Electrostatic stabilization or shielding of negative charges** as Mg²⁺ binding to ATP

ionization limited
should sound

6- Electrostatic catalysis

- Enzymes seem ^(تظهر) to arrange **active site charge distributions** to stabilize the **transition states** ^{حالات انتقالية} of catalyzed reactions
- Substrate binding generally excludes water from an enzyme active site generating a **low dielectric constant** within the active site
- Electrostatic interactions are stronger
- pka's can vary by several pH units due to proximity of charged groups

^{الشكل المخالف}
- Alternative form of electrostatic catalysis: several enzymes as **superoxide dismutase** apparently **use charge distributions** to guide **polar substrates** to their active sites

↓ (SOD)

تستخدم
على

Enzyme Specificity

- In general, there are four distinct types of specificity:

- 1- Absolute specificity: the enzyme will catalyze only one reaction.
- 2- Group specificity: the enzyme will act only on molecules that have specific functional groups, such as amino, phosphate and methyl groups
- 3- Linkage specificity: the enzyme will act on a particular type of chemical bond regardless of the rest of the molecular structure
- 4- Stereo chemical specificity: the enzyme will act on a particular steric or optical isomer.

not for All enzymes (require cofactors)

- Some enzymes require cofactors to be active.

- Cofactors are a non-protein components of the enzyme.


- Organic Molecules (Coenzymes) ✓

- Inorganic ions e.g., Ca^{2+} , Zn^{2+} (Prosthetic group) ✓


- Cofactors may be:

1- The Permanently attached cofactors, are called Prosthetic group (such as a vitamin, sugar, or lipid or inorganic such as a metal ion)


2- Temporarily attached cofactors are called coenzyme, its detach after a reaction and may participate in the reaction with other enzyme.



Enzyme is protein only
Example: lysozyme



Enzyme + prosthetic group
Example:
flavoprotein + FAD



Enzyme + coenzyme
Example:
dehydrogenases + NAD

Cofactors

- Cofactors can be subdivided into two groups: metals and small organic molecules
- Cofactors that are small organic molecules are called coenzymes.
- Most common cofactor are also metal ions.

- If tightly bound, the cofactors are called prosthetic groups.

non-covalent and covalent bond

- Loosely bound Cofactors serve functions similar to those of prosthetic groups but bind in a transient, dissociable manner either to the enzyme or to a substrate

work in more than one of enzymes.

NAD^+ / B_3 } \Rightarrow is a coenzyme

Prosthetic groups

- {Tightly integrated} into the enzyme structure by covalent or non-covalent forces. e.g.;

Ⓐ Pyridoxal phosphate *from vitamin (B6)*

Flavin mononucleotide (FMN) *from (B2)*

Flavin adenine dinucleotide (FAD) *3*

Thiamin pyrophosphate (TPP) *from (B1)*

Biotin \Rightarrow *B7 (its existing as prosthetic group only), (not as coenzyme)*

Ⓑ Metal ions – Co, Cu, Mg, Mn, Zn

- Metals are the most common prosthetic groups

Coenzymes

- Very often vitamins / (also have metal ions)
- They serve as recyclable shuttles—or group transfer agents—that transport many substrates from their point of generation to their point of utilization.
- The water-soluble B vitamins supply important components of numerous coenzymes.
- Chemical moieties transported by coenzymes include hydrogen atoms or hydride^{H⁻} ions, methyl groups (folates), acyl groups (coenzyme A), and oligosaccharides (dolichol).

مرافقة الإنزيم (إنزيم)

Important Prosthetic Groups and Coenzymes

Prosthetic Group	Enzymes/ Proteins
Zn ⁺⁺	Carbonic anhydrase , Alcohol dehydrogenase
Fe ⁺⁺⁺ or Fe ⁺⁺	Hemoglobin, Cytochromes, ferredoxin
Cu ⁺⁺ or Cu ⁺⁺⁺	Cytochrome oxidase
K ⁺ and Mg ⁺⁺	Pyruvate Phosphokinase

Coenzymes	Vitamins
Nicotinamide adenine dinucleotide (NAD ⁺) or nicotinamide adenine dinucleotide phosphate (NADP ⁺)	vitamin B ₃ (niacin)
Flavin mononucleotide (FMN ⁺) or flavin adenine dinucleotide(FAD ⁺)	vitamin B ₂ (riboflavin)
Pyridoxal phosphate	vitamin B ₆ (pyridoxine)
Coenzyme A ✗	Pantothenic Acid

Diagnostic significance of enzymes


- 1- Enzymes can act as diagnostic markers of underlying diseases .
- 2- Enzymes can also act as reagents for various biochemical estimations and detections \rightarrow *glucose oxidase reaction estimation the concentration in the body.* } *the normal sugar in the body is 80-100.*

Enzymes as diagnostic markers

- 1- Functional plasma enzymes ^{*in the blood*} (Plasma derived enzymes):
 - Certain enzymes, proenzymes, and their substrates are present at all times in the circulation of normal individuals and perform a physiologic function in the blood. } *The problem when it found in the cells*

Examples of these functional plasma enzymes include lipoprotein lipase, pseudo cholinesterase, and the proenzymes of blood coagulation and blood clot dissolution .
The majority of them are synthesized in and secreted by liver.

- ## 2- Nonfunctional plasma enzymes (Cell derived enzymes):
- Plasma also contains numerous other enzymes that perform no known physiologic function in blood.
 - These apparently nonfunctional plasma enzymes arise from the routine normal destruction of erythrocytes, leukocytes, and other cells. *the problem when it found in the plasma.*
 - Tissue damage or necrosis resulting from injury or disease is generally accompanied by increases in the levels of several nonfunctional plasma enzymes.



Isoenzymes (Isoenzymes) (A)

- Are homologous enzymes that catalyze the same reaction but have differences in enzymatic properties.
- Often different isoenzymes are found in different locations in a cell or in different organs/tissues of an organism.
- They are from different polypeptide chains that coded by different genes and so, they are affected by different activators and different inhibitors in different tissues.

e.g.:

Lactate dehydrogenase isoenzymes, 4 Polypeptide chains.

- The enzyme interconverts lactate and pyruvate (LDH)
- Humans have two isoenzymic chains for lactate dehydrogenase: LDH (M) found in muscle and LDH (H) found in heart.
- M is optimized to work under anaerobic conditions and H optimized to work under aerobic conditions.

— oxidation reaction —

- There are 5 different isoenzymes.
- The relative ratio of the isoenzymes depends on the location in the organism as well as the developmental stage.

Isoenzyme	Tissue origin
LDH1 (H4) <i>also used for diagnosis the myocardial infarction.</i> <u>130 kDalton</u>	(Cardiac and kidney)
LDH2 (H3M)	Cardiac, kidney, brain and RBCs
LDH3 (H2M2)	Brain, lung and WBCs
LDH4 (HM3)	Lung, skeletal muscle
LDH5 (M4) <i>the havey stone</i>	Skeletal muscle and liver

CK/CPK Isoenzymes

*استخدم اكثر من
Bia marker*

- There are three Isoenzymes.
- Measuring them is of value in the presence of elevated levels of CK or CPK to determine the source of the elevation.
- Each isoenzyme is a dimer composed of two protomers 'M' (for muscles) and 'B' (for Brain).
- These isoenzymes can be separated by, electrophoresis or by ion exchange chromatography.

Isoenzyme	Electrophoretic mobility	Tissue of origin	Mean % in blood
<u>MM</u> (CK3)	Least	Skeletal muscle Heart muscle	97-100%
<u>MB</u> (CK2)	Intermediate <i>most one to diagnose the myocardial infarction</i>	Heart muscle	0-3%
<u>BB</u> (CK1)	Maximum	Brain	0%

Enzyme Kinetics

- It is the field of biochemistry concerned with the quantitative measurement of the rates of enzyme-catalyzed reactions and the study of the factors affecting these rates.
- The rate of a chemical reaction is described by the number of molecules of reactant(s) to be converted into product(s) in a specified time period which is dependent on the concentration of the chemicals involved in the process and on rate constants that are characteristic of the reaction.