Which one of the following measures considered as a primary prevention for Musculo-skeletal injurised one: a. Motorcycle helmets b. life vests c. Trigger locks on guns d. Bullet proof vests e. Seatbelts	ry?

51) A bodily lesion at the organic level, resulting from acute exposure to energy in amounts that exceed the threshold of physiological tolerance called?

- a. Mortality
- b. Disaster
- c. Distress
- d. Morbidity
- e. Injury

Ouestion 17

Not yet answered

Marked out of 1.00

P Flag question

O a. Motorcycle helmets

C. Trigger locks on guns

O d. Bullet proof vests

O e. Seatbelts

18. the best outcome measure of musculoskeletal injuries is:

A disability adjusted life years

- в. mortality adjusted life years
- c. morbidity adjusted life years
- D. injury adjusted life years
- E. morbidity and mortality rate
- 19 If the fatality rate due to deadly musculoskeletal

61) Which one of the following injuries resulted from an insufficiency of a vital element?

- A. Chemical
- B. Electrical
- C. Thermal
- D. Strangulation —
- E. Mechanical

31) Which one of the following measures considered as a primary prevention for Musculo-skeletal injury?

- A. Bullet proof vests
- B. Life vests
- C. Seat belts
- D. Motorcycle helmets
- E. Safety caps on poisonous substances

The single leading cause of disability globally is?

- a. Low back pain
- O b. Neck stiffness
- O c. Ligament strain
- O d. Muscle rupture
- O e. Vertebral block

20. fatal and non-fatal musculoskeletal injuries are more common in:

port places

- н. schools
- 1. homes and around homes
- J. shopping centers

All the	e following statements are considered as unintentional injuries EXCEPT?
Selec	ct one:
0	a. Exposure to venomous plants and animals
0	b. Exposure to animate and inanimate mechanical forces
•	c. Homicide
0	d. Road traffic injuries
0	e. Exposure to electric current, radiation and extreme ambient temperature