


# Cell Bio

## Introduction to stem cell



**Stem cells have the potential to change life as we know it**

# What is a stem cell?

*“Any time you have healed after an injury it’s a stem cell mediated event”*

## unprogrammed cells that are:

- Capable of dividing and renewing themselves for long periods of time i.e. Proliferation
- Have the potential to give rise to specialized cell types i.e. Differentiate
- Stem cell is **unique** because it Can do both:


**Self-renew**

Make copies of itself

**Differentiate**

Make other types of cells  
(specialized cells of the body)


# What is a stem cell?



# Why self renew & Why differentiate?

## A- Self renewal

- Because if they didn't copy themselves, they would **finish** quickly.
- It is important for the body to **maintain a storage** of stem cells to use throughout your life.



## B- Differentiation

- Specialized cells are mature cells **cannot divide** or make copies of themselves, so if they damage or die they need to be replaced so the body can keep on working.
- Specialized* or *differentiated* cells have **particular roles** in the body e.g. blood cells, nerve cells, muscle cells

- There are 2 theories for stem cell division:


**1- Obligate asymmetric stem cell replication**

**2- Stochastic differentiation**

**progenitor (intermediate) cells**: Cells that are at a stage between stem cells and mature specialized cells.


they are yet **not fully programmed**, somehow more mature than a stem cells.

They have less capacity to self-renew themselves than a stem cells



**Potency**: stem Cell's ability to differentiate into other cell types

# 1- Obligate asymmetric stem cell replication



**1 stem cell**




**Self renewal - maintains the stem cell pool**

**4 specialized cells**



**Differentiation - replaces dead or damaged cells throughout your life**

## 2- Stochastic differentiation



If one stem cell differentiate into 2 specialized cells another stem cell will notice that and make up for the lost stem cell and divide by mitosis to produce 2 identical stem cells

# Types of stem cells

There are 2 types of stem cells:

1. Embryonic stem cells (ES)
2. Tissue (adult /somatic) stem cells (TS)

Embryonic stem cells:


Found in **inner cell mass of blastocyst** (a very early stage of embryo life that has about 50 to 100 cells)

Tissue (adult /somatic)stem cells:


found in the **tissues of the body** (in a fetus, baby, child or adult).

# 1- Embryonic stem cells (ES)

- Embryonic stem (ES) cells derived from the inner cell mass of a blastocyst (an early- stage embryo)
- Human embryos reach the blastocyst stage 4-5 days post fertilization (consists of **50–150 cells**) it is the stage at which implantation occurs
- ES of the inner cell mass are **pluripotent**.  
*(Pluripotent: can give rise to all of the cell types that make the body)*
- Few weeks later the cells will organize into 3 primary cell layers → germinal layers : (**ectoderm, mesoderm, endoderm**) **No more pluripotent**. As development continue the cells of that layers will differentiate to form > 200 types of cells that form the body



# Embryonic stem cell



ES cells don't contribute to the extra-embryonic membranes or placenta

*The 3 primary cell layers are formed in the earliest stages of the embryonic development*

The cells in each germ layer differentiate into tissues and organs


- **Ectoderm** → skin, nervous system, & parts of head & neck
- **Mesoderm** → muscles, blood, blood vessels, & beginning of bone & connective tissue
- **Endoderm** → digestive, respiratory tracts, pancreas & liver

### Embryonic stem cell research

**ES cells** offer hope for new therapies, but their use in research has been **strongly debated** because:

1. Destruction of embryo
2. Rejection due to different genetic background
3. Changed into tumor cells. Once they put in the body they can never be taken out






# Embryonic stem cells



ECTODERM

MESODERM

ENDODERM



Skin

Nerves

Bones


Muscles

Liver

Thyroid

- Stem cells is surrounded by a special microenvironment **called the stem cell niche**. Niches consist of a multiple factors that can influence stem cell behavior.
- Any human ES cell is defined by the expression of **several transcription factors on its cell surface**


- The transcription factors **Oct-4, Nanog, Sox 2, max, Smad 1, FoxC2**  
*(Proteins that control rate of Transcription of genetic formation)*



- These factors control the expression of **genes** that either maintain ES pluripotency or induce ES differentiation into progenitors of 3 germ layers


# Signals to stem cell

Niches




## 2- Tissue (adult/somatic ) stem cells

- Undifferentiated cells, found among differentiated cells in tissues & organs **after birth .. Used as repair system**
- They are small in number, have restricted ability to self- renew itself & to differentiate into various types of cells
- Its origin in mature tissue is unknown
- They **differentiate** only to specialized cells similar that of the tissue in which they are found
- They are **multipotent**




- **Therapy using Adult stem cells** has been successfully used for many years to treat **leukemia** and related bone/blood cancers through bone marrow transplants.



- The use of adult stem cells in research & therapy **is not as controversial as** the use of ES cells, because the production of adult stem cells does not require the destruction of an embryo

## Where are adult stem cells found?




- Adult stem cells exist throughout the body from the time an embryo develops
- Adult stem cells replace cells that are damaged or used up.

## The bone marrow contains at least two kinds of stem cells:

A- One is called hematopoietic stem cells, forms all the types of blood cells in the body.

B- The second is called Mesenchymal stem cells (bone marrow stromal cells) that generates bone, cartilage, fat, and fibrous connective tissue cells.

- Stem cells are thought to reside in a specific areas of each tissue where they may remain quiescent (non-dividing) for many years until they are activated by **signals** e.g. disease or tissue injury.



## Cord blood stem cells

- Umbilical cord blood was once discarded as waste material but is now known to be a useful source of blood stem cells.
- After a baby is born, cord blood in the umbilical cord & placenta is relatively easy to collect it contains Hematopoietic (blood) stem cells (will give rise to red cells, white cells, platelets, & lymphocytes)
- is used to reconstitute bone marrow following radiation treatment for various blood cancers, and for various forms of anemia



# Induced pluripotent stem cells (iPSC)

- iPSC technology is a huge discovery (2006) → NP 2012
- Concept: mature cells can be reprogrammed to become pluripotent.
- Technique : done by introduce a few specific pluripotency genes into already specialized cells ( Ex: ms cells) → the cells will forget what type of cells they are & revert back & reprogrammed back into pluripotent stem cell
- Goal: regenerative medicine ..... To replace damage tissue in a given person by using pluripotent stem cells from his own body, not only the patient will get the new organ he needs but also NO any immune- rejection complication


# Induced pluripotent stem cells (iPSC)

1<sup>st</sup> isolate and culture skin cells from a patient.

2<sup>nd</sup> introduce three or four pluripotency genes into the skin cells by using an engineered virus carrier.

The expression of these genes regenerates the stem cell phenotype.


The viruses simply deliver the genes of interest and are themselves engineered not to be harmful.



# Committed progenitor cells

- They are early descendants of stem cells but they are more specific than stem cells

- They have the tendency to differentiate into specific type of cells.




- The most important difference between **stem cells** and **progenitor cells** is that:

**stem cells** can replicate indefinitely, whereas

**progenitor cells** can divide only for limited number of times.

# Committed progenitor cells



## stem cell:

- self renew, divide
- high potency



## committed progenitors:

- "Temporary amplifying cells"
  - multipotent
  - Divide rapidly
  - No self-renewal

## specialized cells

- Function
- No division

- **Genetic & environmental factors** determine the pathway of differentiation that the progenitor cells will take to form a specific lineage . They remain **dormant** in the tissue till need.
- Adult stem cells main role is to replace cells in case of tissue injury, damage or dead cells
- Example of adult stem cells:
  - ✓ **Satellite cells** found in muscle → myogenic progenitor cells → skeletal muscle cells
  - ✓ **Periosteum** contains progenitor cells that develop into osteoblasts.




**Potency:**


Cell's ability to differentiate into other cell types

**Totipotent:**

can give rise to an entire functional organism  
{cells from early embryo (1-3 days)}



**Pluripotent:** can give rise to **all** types of specialized cells in the body (ESC: 5- 14 days)  
(form > 200 cell types)




**Multipotent:** can give rise to **multiple** types of specialized cells, but not all cell types  
(Adult SC in tissues & hematopoietic SC in cord blood)

**Oligopotent:** can differentiate into few cell types  
e.g. lymphoid or myeloid stem cells


**Unipotent:** can give rise to **only one** type of cells.  
e.g. B-lymphocyte → plasma cells  
Monocyte → macrophages

**Nullipotent:** Terminal cells



# Myeloid and lymphoid stem cells






# Embryonic stem cells



ECTODERM

MESODERM

ENDODERM



Skin

Nerves

Bones

Muscles

Liver

Thyroid

# Cloning



- The process of producing a population of genetically identical individuals ( **exact genetic copies** )

- There are 2 types:

## Reproductive cloning:

- Hit the headlines in the late 1990s when '**Dolly the sheep**' was cloned.
- It was the first mammal ever to be cloned.




## Molecular cloning:

- a technique used to help scientists investigate what particular genes do and how they work.

# Cloning


There are two VERY different types of cloning:

## Reproductive cloning



- Use to make two identical individuals
- Very difficult to do
- Illegal to do on humans

## Molecular cloning



- Use to study what a gene does
- Routine in the biology labs

# Reproductive cloning

To make Dolly, scientists done what so called somatic cell nuclear transfer (SCNT)

**Somatic cell**: somatic cell is any cell in the body other than sperm & egg. Somatic cell has the 2 complete sets of chromosomes (46 = 23pairs)

**Nuclear** : nucleus holds DNA which contains all the information needed to form an organism


**Transfer**: moving an object from one place to another

## Technique

- They took the nucleus out of a normal somatic cell from a sheep (original).
- They put that nucleus into an egg cell of another sheep that had no nucleus.
- They then had a new cell.
- To make the new cell start to divide and grow, they gave it an electric shock.

- Then it started to divide and develop into an embryo. When it had grown into a very early stage embryo called a blastocyst
- it then was implanted into the womb of another sheep so that it could grow into a lamb and be born.
- The new sheep is a clone of the sheep that donated the somatic cell. Both sheep have the same DNA

# Reproductive cloning




# Molecular cloning

- A process used by scientists to make **copies of a specific gene or genes** inside a cell, in order to study what these genes do or how they work?


- Also called **recombinant DNA cloning**

- Recombinant DNA:


- DNA molecules formed by laboratory methods of genetic recombination to bring together genetic material from multiple sources → creating DNA sequences that would not found in the genome



- Recombinant DNA is possible because DNA molecules from all organisms share the same chemical structure ( the genetic code is universal) .
- Construction of recombinant DNA, involves insertion of a foreign DNA fragment into a plasmid vector.




- **Cloning vector:** It carries the genetic material of interest into another cell where it can be replicated &/or expressed
- A small piece of DNA taken from a **virus** or a **plasmid**, into which a foreign DNA fragment can be inserted and can be stably maintained in an organism for cloning purposes



- **Plasmid:** circular double stranded DNA molecule within a bacterial cell (physically separated from chromosomal DNA) can replicate independently

- **virus** is a small infectious agent that replicates only inside the living cells of other organisms. Viruses can infect all types of life forms.



## Technique


- 1- Select the **DNA molecule to be cloned** (of interest)
- 2- Select **DNA molecule** that will serve as a **vector (virus /plasmid)**
- 3- cleave the vector DNA strand with **Restriction endonuclease**, then insert foreign DNA → recombinant DNA
- 4- Introduction of recombinant DNA in host cells → transformation
- 5- When the cell divides, it makes copies of itself. Each new daughter cell contains an exact copy of the new DNA (cloned DNA)
- 6- Selection & screening of colonies with desired DNA

# Applications of molecular cloning:

**1- Insulin production:** we use the bacteria to be human insulin factories

Why bacteria?

- Contain plasmids
- Are unicellular and reproduce asexually → quick clones



## 2- Study genome organization and gene expression

### A- Loss of function (gene knockout):

A genetic technique in which a gene is removed or blocked so that *it does not work*, used in learning about a gene that has been sequenced but has an unknown or incompletely known function

B- Transgenic organisms :generating genetically modified organisms (GMOs), most GMOs are generated for purposes of basic biological research e.g. transgenic mouse

C- Gene therapy: involves supplying a functional gene to cells lacking that function, with the aim of correcting a genetic disorder or acquired disease

# Thank you

