يمنع أخذ السلايدات بدون إذن المحرر واي اجراء يخالف ذلك يقع تحت طائلة المسؤولية القانونية جميع المعلومات للاستخدام التعليمي فقط

Juistral 3. 22 Apple 201




الأستاذ الدكتور يوسف حسين

كلبة الطب - جامعة مؤتة - الأردن

ة من جامعة كولونيا

Radioulnar Joints



Superior Radioulnar Joint

** Type; a pivot (uniaxial) synovial joint.

** Articular surfaces:

- Articular circumference of the head of the radius.
- 2- Radial notch of the ulna and annular ligament.

Annular ligament:

- It is a strong fibrous band surrounding the head of the radius.
- It is attached to the margins of radial notch of the ulna.

Radius Ulna Synovial membrane: lines inner surface of capsule and Continuous with membrane of elbow joint.


of ulna

Radial head

Annular ligament

Inferior Radioulnar Joint

- ** Type; a pivot (uniaxial) synovial joint.
- ** Articular surfaces;
- Articular circumference of the head of ulna
- Ulnar notch of radius.
- ** Articular disc; is a triangular fibrocartilage.
 - Its apex attached to base of styloid process of ulna.
 - The base is attached the inferior margin of the ulnar notch of the radius.
 - The disc separates the inferior radioulnar joint above from wrist joint below.
 - ** Synovial membrane, projects upwards forming recessus sacciformis in front the interosseus membrane.




 Movements of superior and inferior radioulnar joints

Palm anterior

- Pronation and Supination
- Definition:
 - Pronation: medial rotation of the forearm and the palm of the hand faces posteriorly.
 - Supination: lateral rotation of the forearm and the palm of the hand faces anteriorly.
- inferior Joints: superior and radioulnar joints.






Supination

- a- It is initiated by brachioradialis.
- b- It is completed by
- Supinator during extension of elbow joint.
- Biceps brachii during flexion of elbow joint.
- Radius is parallel to ulna
- N.B Supination is more powerful than pronation. Because, in usual working position with elbow flexed (e.g. Turning screwdriver).
- In a semiflexed elbow: palm is turned upwards in supination and downward in pronation (king pronates and beggar supinates)

Supination





Radioulnar Joints

- Middle Radioulnar Joint
- ** Type; fibrous joint (No movements). It is formed by
- 1- Oblique cord: is a fibrous band extending from ulnar tuberosity to shaft of radius below the radial tuberosity.
- 2- Interosseous membrane: thin sheet of fibrous tissue between interosseous borders of the radius and ulna.
- Directions of fibres, obliquely downwards and medially from the radius to the ulna.
- Functions of interosseous membrane:
 - 1- It binds radius and ulna.
 - 2- Breaks the shock transmitted from hand to radius.
 - 3- It gives origin for the deep muscles of the forearm.
- Openings,
 - 1- Anterior interosseous artery perforates membrane to the back of the forearm above the upper border of pronator quadratus.
 - 2- Posterior interosseous artery passes in gap between the oblique cord and interosseous membrane to the back of the forearm.

 Prof. Dr. Youssef Hussein Anatomy YouTube


Oblique cord Opening of posterior interosseus artery Ulna

Interosseous membrane

Opening of anterior interosseus artery

Inferior radio-ulnar joint
Articular disc





Palmar aponeurosis

- It is a thickening of the deep fascia in the middle of the palm
- It improve the hand grip
- It protects the underlying tendons, blood vessels and nerves

** Shape; - It is a triangular.

The apex is adherent to the flexor retinaculum and gives insertion to the tendon of palmarislongus.

 The base divides into four slips, one for each of the medial four fingers.


** Dupuytren's contracture

Thickening and fibrosis of the palmar aponeurosis due to its inflammation leading permeant flexion to of metacarpophalangeal and interphalangeal proximal joints of the 4th (ring) 5th (little finger)fingers. It is painless condition










4- Adductor pollicis


3- Opponens Pollicis deep to abductor and flexor pollicis brevis


2- Flexor pollicis brevis

1-Abductor pollicis brevis

- Nr. 1,2,&3 called thenar eminence
- Nerve supply: All thenar muscles supplied by lateral division of median nerve except adductor pollicis by deep branch of ulnar nerve.
- Thenar eminence has the same nerve & origin









4- Adductor Pollicis

** Origin: by two heads:

a- Oblique head: from the bases of the 2nd and 3rd metacarpal and carpal bones.

b- Transverse head: from the palmar surface of the 3rd metacarpal bone.

* Insertion: into the medial side of the base of the proximal phalanx of the thumb.


* Nerve supply: deep branch of the ulnar nerve.


* Actions: adduction of the thumb.


N.B; A small sesamoid bone is developed in the tendon of adductor pollicis.














Hook of hamate

3- Opponens Digiti Minimi

** Site; deep to the flexor and abductor digiti minimi

** Origin: a) Medial part of the flexor retinaculum.


b) Hook of hamate.

** Insertion: Into the medial half of the palmar surface of the 5th metacarpal bone.

** Action: opposition of the little finger (It pulls the 5th metacarpal bone slightly forwards and rotates it laterally.


Prof. Dr. Youssef Hussein Anatomy - YouTube

Flexor retinaculum



Lumbricalis muscles Interossei muscles

- All muscles are supplied by deep branch of the ulnar nerve EXCEPT
- The 1st lumbricalis from lateral division of median nerve
- The 2nd lumbricalis from medial division of median nerve



Lumbrical Muscles

 They are 4 muscles arranged from lateral to medial side.


** Origins: tendons of the flexor digitorum profundeus


- 1- The 1st and 2nd are unipennate
- 2- The 3rd and 4th are bipennate.
- ** Insertions extensor expansion of the corresponding finger.
- ** each tendon crosses the lateral side of the corresponding metacarpophalangeal joint.

** Actions:

Flexion of the metacarpophalangeal Joints and extension the interphalangeal joints of the medial 4

fingers (writing position). Prof. Dr. Youssef Hussein Anatomy - YouTube






Palmar interossei

- They are smaller than the dorsal interossei.

** Origin

- 1st from the medial side of the 2nd metacarpal bone.
- 2nd from the lateral side of the 4th metacarpal bone.
- 3rd from the lateral side of the 5th metacarpal bone.







Palmar interossei

** Insertion

- 1st to medial side of the proximal phalanx and extensor expansion of the index.
- 2nd to lateral side of the proximal phalanx and the extensor expansion of the ring.
- 3rd to lateral side of the proximal phalanx and the extensor expansion of the little finger.
- ** Actions: adduct the fingers towards the axis.
- They assist lumbricals in flexion of metacarpophalangeal Joints and extension interphalangeal joints (writing positions).

N.B: No palmar interossei to the middle finger (axis)





Dorsal interossei (4 in number)

- ** Origin from adjacent sides of the metacarpal bones.
- 1- The 1st From the adjacent sides of the 1st and 2nd metacarpal bones.
- (large) and sometimes called abductor indicis.
- 2- The 2nd from the adjacent sides of the 2nd and 3rd metacarpal bones.
- 3- The 3rd from the adjacent sides of the 3rd and 4th metacarpal bones.
- 4- The 4th from the adjacent sides of the 4th and 5th metacarpal bones.



Dorsal interossei are Bipennate

Dorsal interossei


** Insertion:

Prof. Dr. Youssef Hussein Anatomy YouTube


- 1- The 1st into the lateral side base of the proximal phalanx and extensor expansion of the index.
- 2- The 2nd into the lateral side of the base of the proximal phalanx and extensor expansion of the middle finger.
- 3- The 3rd into the medial side of the base of the proximal phalanx and extensor expansion of the middle finger.
- 4- The 4th into the medial side of the base of the proximal phalanx and extensor expansion of the ring finger.
- ** Actions: abduct the fingers away from the axis of the middle linger.
- They assist lumbricals in flexion of metacarpophalangeal
 Joints and extension interphalangeal joints (writing positions).

N.B: No dorsal interossei to thumb and little fingers

- The axis receives two (2nd & 3rd) dorsal interossei








Fascial Spaces In the hand

- Two palmar septa (medial and lateral) From palmar aponeurosis divide palm into 3 main fascial compartments.
- I- Lateral compartment: contains muscles of thenar eminence.
- 2- Medial compartment: contains muscles of hypothenar eminence.
- 3- Intermediate compartment: is the wide space deep to the palmar aponeurosis.
- An intermediate septum extends from the deep surface of palmar aponeurosis between flexor tendons of the index laterally and the 2nd lumbrical medially.
- It divides this compartment into medial (midpalmar space) and lateral (thenar space) parts.



Prof. Dr. Youssef Hussein Anatomy - You Tube

a- Thenar space

** Contents;

a- Flexor tendons of the index.

b- The first lumbricals. Prof. Dr. Youssef Hussein Anatomy - You Tube


c- Digital vessels and nerves of the lateral one and half fingers.

N.B. surgical anatomy: It lies superficial to adductor pollicis forming a deep connection between radial and ulnar bursa leading to spread of infection in the palm of the hand


b- Midpalmar space

Contents;

- a Flexor tendons of the middle, ring and little fingers.
- b- The 2nd, 3rd and 4th lumbricals
- c- Digital vessels and nerves of the medial three and half fingers.
- D- Superficial and deep palmar arches and their branches



Digital Pulp Space



Pulp space

- They are in the palmar surface of the tips of the fingers distal to the insertion of the flexor digitorum profundus.
- They contain fatty tissue that is divided into numerous compartments by fibrous septa.
 Prof. Dr. Youssef Hussein Anatomy YouTube
- The terminal branches of the digital vessels pass through the pulp.
- It is closed proximally by the fusion of the deep fascia to the periosteum of the base of the terminal phalanges.

* Clinical importance;

- 1- Pulp space infection (whitlow) produce severe throbbing pain due to closed space.
- 2- Infection of this space is drained by a lateral incision
- 3- Severe infection of the pulp space may lead to occlude the digital arteries with consequent necrosis of the terminal phalanx.

Web spaces

 Spaces between the bases of the proximal phalanges of the fingers containing connective tissue and fat.



The 1st web space

Prof. Dr. Youssef Hussein Anatomy - YouTube



Prof. Dr. Youssef Hussein Anatomy - YouTube



Power grip

Holding a heavy object firmly between the fingers thumb.

Prof. Dr. Youssef Hussein Anatomy - YouTube

Precision grip

Holding a small object lightly and precisely between the tip of the thumb and the tip of the index with partial support by the middle finger e.g. holding of the pencil in the writing position or surgical scalpel.





Hook grip

As for carrying a handle bag, done by the long flexor tendons.

Branches of radial artery

Branches of ulnar artery

- A- in the forearm
- 1- Radial recurrent artery
- 2- Palmar carpal branch,

Arch

Arch

Arch

- 3- Superficial palmar branch
- B- in the dorsum of the hand
- 1- Dorsal carpal branch -
- 2- Dorsal digital branch
- 3- First dorsal metacarpal artery
- C- in the palm of the hand
- 1- Princeps pollicis artery
- 2- Radialis indicis artery
- D- End of the radial artery

- A- in the forearm
- I- Anterior ulnar recurrent artery
- 2- Posterior ulnar recurrent artery
- 3- Common interosseous artery
- 5- Palmer carpal branch
- 6- Dorsal carpal branch


- B- in the palm of the hand
- 1- Superficial branch
- 2-Deep branch




Dorsal carpal arch

Dorsal Carpal Arch

- * **Formation**; union of:
- 1- Dorsal carpal branch of radial artery.
- 2- Dorsal carpal branch of ulnar artery.
- 3- Termination of anterior interosseous artery.
- * Branches
- 1- 2nd, 3rd and 4th dorsal metacarpal arteries to the adjacent sides of the medial 4 fingers.
- 2- Dorsal digital artery for the medial side of little finger.








Superficial palmar arch

- ** Site: superficial to the flexor tendons.
- It is distal to the deep palmar arch.
- ** Formation;
- 1. Superficial terminal branch of the ulnar artery (mainly).
- 2. Superficial palmar branch of the radial artery.
- ** Branches:
 - 1- 3 Palmar digital arteries for the adjacent sides of medial 4 fingers
 - 2- Palmar digital artery for the medial side of the little finger.
- Pulp Spaces are fibrofatty compartments on the palmar surface of the distal phalanges distal to the insertion of the flexor tendons. They contain ramification of the digital arteries (artery of pulp). Infection of the pulp space is painful.
 Obstruction of these arteries leading to necrosis.







Deep palmar arch

** Site; deep to the flexor tendons and lumbrical muscles.

** Formation;

- 1- The continuation of the radial artery (mainly).
- 2- The deep terminal branch of the ulnar artery.

** Branches

1-3 palmar metacarpal arteries join palmar digital arteries of superficial palmar arch.

YOUTUBE. يوسف حسين

- 2- 3 Perforating branches: pass through the 2nd, 3rd and 4th interosseous spaces to join the 2nd, 3rd and 4th dorsal metacarpal arteries of dorsal carpal arch.
- 3- Recurrent branches to join the palmar carpal arch.
- In cases of hemorrhage of the hand press the radial artery against lower end of the radius to identify the source of bleeding;
- a- If it becomes weak, the source of bleeding is the deep palmar arch.
- b- If not changed, the source of bleeding is the superficial palmar arch.

https://www.youtube.com/channel/UCVSNqbibj9UWYaJdd_cn0PQ



https://www.youtube.com/@ProfDrYoussefHusseinAnatomy/playlists